

UNCLASSIFIED

AD NUMBER
ADB244278
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies only; Proprietary Info.; Aug 98. Other requests shall be referred to U.S. Army Medical Research and Materiel Comd., 504 Scott St., Fort Detrick, MD 21702-5012.
AUTHORITY
USAMRMC ltr, 23 Aug 2001

THIS PAGE IS UNCLASSIFIED

AD_____

AWARD NUMBER DAMD17-97-1-7241

TITLE: A Novel Repressor of Estrogen-Regulated Genes for Breast Cancer Growth Suppression

PRINCIPAL INVESTIGATOR: David J. Shapiro, Ph.D.

CONTRACTING ORGANIZATION: University of Illinois
Champaign, Illinois 61820

REPORT DATE: August 1998

TYPE OF REPORT: Annual

PREPARED FOR: Commander
U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Distribution authorized to U.S. Government agencies only (proprietary information, Aug 98). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

DTIC QUALITY INSPECTED 4

NOTICE

USING GOVERNMENT DRAWINGS, SPECIFICATIONS, OR OTHER DATA INCLUDED IN THIS DOCUMENT FOR ANY PURPOSE OTHER THAN GOVERNMENT PROCUREMENT DOES NOT IN ANY WAY OBLIGATE THE U.S. GOVERNMENT. THE FACT THAT THE GOVERNMENT FORMULATED OR SUPPLIED THE DRAWINGS, SPECIFICATIONS, OR OTHER DATA DOES NOT LICENSE THE HOLDER OR ANY OTHER PERSON OR CORPORATION; OR CONVEY ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE, OR SELL ANY PATENTED INVENTION THAT MAY RELATE TO THEM.

LIMITED RIGHTS LEGEND

Contract Number: DAMD17-97-1-7241

Contractor: University of Illinois

Location of Limited Rights Data (Pages): 4-28

Those portions of the technical data contained in this report marked as limited rights data shall not, without the written permission of the above contractor, be (a) released or disclosed outside the government, (b) used by the Government for manufacture or, in the case of computer software documentation, for preparing the same or similar computer software, or (c) used by a party other than the Government, except that the Government may release or disclose technical data to persons outside the Government, or permit the use of technical data by such persons, if (i) such release, disclosure, or use is necessary for emergency repair or overhaul or (ii) is a release or disclosure of technical data (other than detailed manufacturing or process data) to, or use of such data by, a foreign government that is in the interest of the Government and is required for evaluational or informational purposes, provided in either case that such release, disclosure or use is made subject to a prohibition that the person to whom the data is released or disclosed may not further use, release or disclose such data, and the contractor or subcontractor or subcontractor asserting the restriction is notified of such release, disclosure or use. This legend, together with the indications of the portions of this data which are subject to such limitations, shall be included on any reproduction hereof which includes any part of the portions subject to such limitations.

THIS TECHNICAL REPORT HAS BEEN REVIEWED AND IS APPROVED FOR PUBLICATION.

James C. Morrow 5/20/99

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.</small>				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE August 1998		3. REPORT TYPE AND DATES COVERED Annual (1 Aug 97 - 31 Jul 98)
4. TITLE AND SUBTITLE A Novel Repressor of Estrogen-Regulated Genes for Breast Cancer Growth Suppression			5. FUNDING NUMBERS DAMD17-97-1-7241	
6. AUTHOR(S) David J. Shapiro, Ph.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Illinois Champaign, Illinois 61820			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research And Materiel Command ATTN: MCMR-RMI-S 504 Scott Street Fort Detrick, Maryland 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES			<div style="border: 1px solid black; padding: 10px; text-align: center; font-size: 2em; font-weight: bold;">19990603 074</div>	
12a. DISTRIBUTION / AVAILABILITY STATEMENT Distribution authorized to U.S. Government agencies only (proprietary information, Aug 98). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.				
13. ABSTRACT (Maximum 200 words) <p>The objective of this project is to develop and use novel ER mutants to test a two-part hypothesis. First, that the estrogen-independent growth of breast cancer cells involves the estrogen-independent expression of growth control genes which are normally estrogen regulated. Second, that suppressing the expression of these genes will block the growth of breast cancer cells. To test these ideas, we set out to develop a novel type of estrogen receptor chimera which will efficiently and quantitatively suppress both estrogen-dependent and estrogen-independent expression of estrogen-regulated growth stimulatory genes. To develop Estrogen Regulated Gene repressors (ERG-repressors), we constructed plasmids encoding a variety of recombinant estrogen receptors fused to different versions of the KRAB repressor domain. Another class of repressors was constructed by fusing the KRAB repressor domain to mutant DNA binding domains selected with the P22 challenge phage system, displaying a strongly enhanced affinity for the estrogen response element (ERE). The activity of these chimeric proteins as ERG-repressors was has evaluated in transient transfection assays and KRAB-estrogen receptor chimeras which are potent gene repressors have been identified.</p>				
14. SUBJECT TERMS Breast Cancer			15. NUMBER OF PAGES 29	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Limited	

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

____ Where copyrighted material is quoted, permission has been obtained to use such material.

____ Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

____ Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.


DS In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

DS For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

DS In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

DS In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

____ In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.


PI - Signature

8/26/98
Date

TABLE OF CONTENTS

FRONT COVER	1
Standard Form (SF) 298, REPORT DOCUMENTATION PAGE	2
FOREWORD	3
TABLE OF CONTENTS	4
INTRODUCTION	5-6
BODY	7
Materials and Experimental Methods	7-11
Results	11-12
Recommendations in Relation to the Statement of Work	13-14
CONCLUSIONS	14
REFERENCES	15
APPENDICES	19-28
Figures	19-27
TABLE	28

INTRODUCTION

Estrogens, acting through the estrogen receptor (ER), play a critical role in regulating the growth and metastases of breast cancers. Growth promotion by estrogen is thought to involve direct estrogen receptor-mediated regulation of the expression of several genes important in cell growth, including those encoding some growth factors (such as TGF- α , IGF1, and their receptors), some early response genes (such as c-myc, and cyclin D1), and other genes, including the progesterone receptor gene. Interference with estrogen activity, usually based on antiestrogens, such as tamoxifen, or on aromatase inhibitors, therefore represents a mainstay in breast cancer treatment. While antiestrogen therapy is often effective initially, the tumors almost always eventually progress to estrogen-independent growth. This limits the long-term utility of endocrine therapies.

It is usually accepted that the ability of the 17 β -estradiol-ER complex to influence the growth and metastasis of breast cancer cells is due to its ability to regulate the expression of specific genes. The estradiol-ER complex acts directly to induce the expression of a set of "early genes" including c-myc, cyclin D1 and TGF- α , and other genes important in cell growth (1-16) by directly interacting with these genes. These early genes, and the products of other directly regulated genes, such as the progesterone receptor, may also initiate a regulatory cascade leading to the regulation of downstream genes important in growth control in breast cancer cells.

It has been widely proposed that the 17 β -estradiol-ER complex induces breast cancer cell growth by directly or indirectly regulating the expression of genes important in cell growth control. **If genes critical to growth control in breast cancer cells are directly induced by the estradiol-ER complex, then repression of estrogen-dependent and estrogen-independent transcription of these genes should block estrogen stimulated growth of the breast cancer cells.** Although it is known that estrogen growth autonomous cells synthesize high levels of growth factors that are normally under estrogen regulation (2-4,6,12,13,17-21), the hypothesis that the high level expression of growth factor genes is responsible for growth of these cells has never been tested directly. We will test this hypothesis by repressing the transcription of these growth factor genes and determining the effect on breast cancer cell growth. If this hypothesis is correct, suppression of the expression of these genes should result in the loss of estrogen-independent growth of these cells, and possibly in their death. In addition, identification of genes as critical for estrogen-dependent or estrogen-independent tumor cell growth provides a basis for the identification of additional agents to suppress their activity.

Background

The KRAB repressor system.

When tethered to a DNA binding domain the KRAB repressor, a 75 amino acid segment found in a number of DNA binding proteins, can efficiently suppress transcription of synthetic genes containing strong binding sites for the protein (22-28). Efficient repression of the natural HIV TAT gene has also been reported (27).

The P22 challenge phage system.

Any sequence-specific DNA binding protein can act as a repressor of transcription if its binding site is close to the transcription initiation site. In our application of the P22 challenge phage system (29, 30), we generated a recombinant phage with an ERE close to the transcription initiation site of the phage P22 Ant gene, whose expression results in lysis of the bacterial host cell. If a steroid receptor mutant with sufficient affinity binds to this ERE it will block the transcription of the gene, allowing the growth of bacterial colonies. Although conceptually quite simple, work with the challenge phage system is quite complex and it required a major development effort to modify it so that it could be used for the first time with a vertebrate protein

Project Overview

We developed a phage selection system which allows us to select estrogen mutants with an enhanced ability to bind to the estrogen response element. The enhanced affinity mutants will therefore bind efficiently to the relatively weak, imperfect, EREs present in most estrogen-regulated genes. In parallel we introduced into various forms of the ER the KRAB box which has been shown to be a potent repressor of transcription of genes to which they are bound. The resulting novel, ligand regulated, ER chimeras are termed estrogen regulated gene-repressors, (ERG-repressors). The ERG-repressors have the capacity to efficiently repress both estrogen-dependent and estrogen-independent transcription from ERE-containing genes in both ER-dependent and ER-independent breast cancer cells. The development of the ERG-repressors, which is detailed below, formed the core objective of our first years work.

In subsequent studies we will express the most effective ERG-repressor in breast cancer cells and determine whether repression of the expression of estrogen-regulated genes blocks the growth of estrogen-dependent breast cancer cell lines, and of our tamoxifen resistant and estrogen (E_2) growth-autonomous lines of MCF-7 cells, and of ER negative MDA-MB-231 cells, both in cell cultures and in tumors formed in immune suppressed mice. These studies will serve as a base for a potential gene therapy approach to breast cancer treatment. In addition these studies will test the hypothesis that expression of ER-regulated genes in breast cancers, is critical to estrogen-independent and estrogen-dependent tumor growth.

BODY

In the first section of the Progress Report which corresponds to Experimental Methods we describe the construction of the numerous estrogen receptor-KRAB chimeras developed and tested in the first year of this project. This section also contains descriptions of other experimental methods used in this work. In the second section of the Progress Report we describe the results obtained when we tested the ability of the candidate ERG-repressors to suppress expression of estrogen-regulated genes.

Materials and Experimental Methods

Construction of ERG-repressors.

p18 KRAB Constructions.

Dr. Luigi Lania (University of Naples 'Frederico II', Italy) provided us with the cDNA of p18KRAB. This protein only contains the KRAB A domain. The KRAB domain consisting of aa 180-235 of the p18 gene was cloned either on the N-terminus or the C-terminus of the human estrogen receptor and a number of transcriptionally inactive dominant negative mutants in the following ways.

P18 KRAB on the N-terminus of hER, hERS554FS and hERL540Q.

The p18 KRAB domain was PCR amplified with the following primers

Forward: TAGAATTCATGCTCCTAACAGCCCAG

Reverse: GAGGGTCATGGTCATGGTCTCCCATTC

The product was then digested with MspI and ligated to the N-terminal MspI/FseI. The resulting ligation product was then digested with EcoRI/NotI. Upon gel purification the resulting fragment was ligated into the pCMV5hER backbone that was ligated with EcoRI/NotI. To obtain the KRAB ER chimeras of the dominant negative mutants hERS554FS and hERL540Q (31, 33), the wild type ligand binding domain was exchanged as an XbaI/BamHI fragment.

KRAB on the C terminus of hER, hERL540Q and Δ A/B-hER

Using the Quikchange mutagenesis strategy (Stratagene) a unique NheI site was introduced at the C-terminal end of the hER and hERL540Q.

Forward: GAGGCAGAGGGTTTCCTGCTAGCTGCCACAGTCTGAG

Reverse: CTCAGACTGTGGCAGCTAGCAGGAAACCCTCTGCCTC

The p18 KRAB domain was PCR amplified using the following oligonucleotides:

Forward: TGAGAGCTAGCAGCCCAGCCCCAGGAG

Reverse: GATCAGCTAGCCCTCGGTCATGGTCTCCCAT

The resulting PCR product was digested with BamHI/XbaI and ligated into the NheI/BamHI digested backbone of hER and hERL540Q. The Ligand binding domain of the resulting ER-KRAB chimeras was obtained as an XbaI/BamHI fragment and ligated

into the backbone of similarly digested $\Delta A/B$ -hER. This same procedure was also followed with wild type ligand binding domain KRAB fusion and the N-terminal KRAB hER backbone to construct an hER-KRAB chimera containing KRAB domains at both the N- and C-terminus.

Cloning of ER KRAB chimeras with ZNF10 (Kox1), ZNF133 and ZNF140 KRAB

The following KRAB domains containing both the KRAB A and KRAB B domain were also used to generate more potent ER KRAB chimeras. The ZNF10 KRAB cDNA (25) was a kind gift of Dr. Hans-Jürgen Thiesen (University of Rostock, Germany), while the ZNF133 and ZNF140 cDNAs (24, 26) were sent to us by Dr. Henrik Vissing (Novo Nordisk, Denmark).

The following fragments from plasmid pCMV5hER were subcloned into vector pGEM11Zf(+) (Promega): 1) the N-terminal fragment EcoRI/NotI; 2) the NotI/HindIII fragment containing the ligand binding domain; 3) The HindIII/BamHI C-terminal fragment of pCMV5hER and pCMV5hERL540Q (31, 33). Quikchange mutagenesis was employed to generate unique NheI sites in these fragments to generate the vectors pG11EnsNhe, pG11EnhNhe, pG11EbhNhe and pG11QbhNhe respectively.

Quikchange primers for pG11EnsNhe

Forward: GCCCGCGGCCACGGACCGCTAGCAATGACCATGACCCTCCA
Reverse: TGGAGGGTCATGGTCATTGCTAGCGGTCCGTGGCCGCGGGC

Quikchange primers for pG11EnhNhe

Forward: AAGTATGGCTATGGAGCTAGCCAAGGAGACTCGCTA
Reverse: TAGCGAGTCTCCTTGGCTAGCTCCATAGCCATACTT

To introduce the NheI site into pG11EbhNhe and pG11QbhNhe the same oligos were used as described above for the cloning of the C-terminal p18KRAB fusion.

Thermocycler sequencing with the BigDye kit (ABI Prism) was employed to verify that the sequences were correct.

For the N-terminal KRAB-ER chimeras the following oligonucleotides were used for PCR amplification using Platinum Taq DNA polymerase(GibcoBRL):

ZNF10 (aa 1-91)

Forward: CAGAATTCATGGATGCTAAGTCACTAAC
Reverse: TATCTAGAAATGCAGTCTCTGAATCAG

ZNF133 (aa 1-119)

Forward: CAGAATTCATGGCATTTCAGGGATGTG
Reverse: TATCTAGAGGCTGGATGTTACCTTCTG

ZNF140 (aa 1-106)

Forward: CAGAATTCATGTAGGGGTCAGTGAC
Reverse: TATCTAGAATTCTTCCATGATCAAATAC

The amplified products were subcloned into the pGEM T-vector (Promega) and the sequence was verified using Thermocycler sequencing with the BigDye kit (ABI Prism). The inserts were obtained as EcoRI/XbaI fragments and together with either the NheI/NotI fragment of plasmid pG11EnsNhe cloned into pCMV5hER digested with EcoRI/NotI or with the NheI/HindIII fragment of pG11EnhNhe into pCMV5hER digested with EcoRI/HindIII. These manipulations generated the following clones:

K10-ER	K10-ΔA/B-ER
K133-ER	K133-ΔA/B-ER
K140-ER	K140-ΔA/B-ER

For the C-terminal KRAB-ER chimeras the following oligonucleotides were used for PCR amplification using PlatinumTaq (GibcoBRL):

ZNF10 (aa 1-91)

Forward:	CTTCTAGATATGGATGCTAAGTCACTAAC
Reverse:	ATGGATCCTAAATGCAGTCTCTGAATCAG

ZNF133 (aa 1-119)

Forward:	CTTCTAGATATGGCATTTCAGGGATGTG
Reverse:	ATGGATCCTAAGGCTGGATGTTACCTTCTG

ZNF140 (aa 1-106)

Forward:	CTTCTAGATATGTAGGGGTCAGTG
Reverse:	ATGGATCCTATCTTTCCATGATCAAATACTG

The amplified products were subcloned into the pGEM T-vector (Promega) and the sequence was verified using Thermocycler sequencing with the BigDye kit (ABI Prism). The inserts were obtained as XbaI/BamHI fragments and ligated into NheI/BamHI digested plasmids pG11EbhNhe and pG11QbhNhe, respectively. The ER LBD-KRAB fusions were obtained as XbaI/BamHI fragments and cloned into pCMV5hER and pCMV5ΔA/B-hER digested with XbaI/BamHI as well as their similarly digested N-terminal chimeras generating the following chimeras:

ERK10	ERQK10
ERK133	ERQK133
ERK140	ERQ140
KERK10	KERQK10
KERK133	KERQK133
KERK140	KERQK140
ΔA/B-ERK10	ΔA/B-ERQK10
ΔA/B-ERK133	ΔA/B-ERQK133
ΔA/B-ERK140	ΔA/B-ERQK140

K-ΔA/B-ERK10
K-ΔA/B-ERK133
K-ΔA/B-ERK140

K-ΔA/B-ERQK10
K-ΔA/B-ERQK133
K-ΔA/B-ERQK140

Flag-GAL4-KRAB constructions

To serve as controls a number of Flag-GAL4-KRAB chimeras were constructed. Dr. Cheng Ming Chiang (University of Illinois) provided us the Flag-GAL4-VP16 fusion protein cloned into the bacterial expression plasmid pET11d (Novagen). For our purpose we sub cloned the coding sequence into a mammalian expression vector for mammalian expression in the following way: the parent plasmid was digested with NcoI and filled in with Pfu polymerase. Subsequently the insert was liberated with a BamHI digest. The gel purified fragment was then ligated into the vector pCDNA 3 (Stratagene) that was digested with HindIII, filled in with Pfu polymerase and subsequently digested with BamHI to generate plasmid pFGVP16. Subsequently the C-terminal end in conjunction with a polylinker was obtained as a PCR fragment from the plasmid pM (Clontech) in which the Dam methylation sensitive BclI site was changed into an ApaI site. The PCR fragment was digested with XhoI/ApaI and ligated into the similarly digested plasmid pFGVP16 to generate plasmid pFGmcs. The N-terminal KRAB domains of ZNF10, ZNF133, ZNF140 were obtained as EcoRI/BamHI fragments and ligated into plasmid pFGmcs, which provided the stop codon generating the vectors pFGK10, pFGK133 and pFGK140, respectively.

Reporter plasmids

To serve as an indicator of repression the following plasmid containing an estrogen responsive promoter displaying high endogenous activity in the absence of estrogen receptor was constructed. The estrogen response elements were obtained from plasmid (ERE)₄-TATA-CAT (33), which was digested with HindIII, blunt ended with Pfu polymerase and religated to generate an NheI site. Subsequently an NheI/BglII digest was performed to liberate the EREs. This fragment was then ligated into the similarly digested vector pGL3-Control (Promega). A similar plasmid was generated for the Flag-GAL4-KRAB constructs in the following way. Five GAL4 binding sites were lifted from the plasmid pG5E1b (34) as an XhoI/BamHI fragment and inserted into either XhoI/BglII or BglII/SalI digested plasmid pGL3-Control to generate plasmids 5'-G5-pGL3-Control and 3'-G5-pGL3-Control, respectively.

Cell maintenance and transfection

The human hepatoma cell line HepG2 and Simian COS-7 cell line were maintained at 37 °C, 5% CO₂ in Dulbecco's Minimal Essential medium (Sigma) supplemented with 10% charcoal dextran stripped fetal bovine serum (Atlanta Biologicals) and 50,000 U/I Penicilline and 50 mg/I Streptomycin (Gibco/BRL).

The hamster cell line CHO was maintained in Dulbecco's Modified Eagle Medium: Nutrient Mixture F-12, 1:1 (Sigma), 29.2 mg/I L-Glutamine (Sigma), 5% charcoal dextran stripped newborn bovine serum (Atlanta Biologicals), 50,000 IU/I Penicilline and 50 mg/I Streptomycin

Cells were transfected with the calcium phosphate coprecipitation method as published previously. Briefly, Cells were plated in 60mm dishes at a density of 4.5×10^5 for HepG2 cells and 2×10^5 for CHO cells, respectively. Forty hours after plating the media was replaced and 2-6 h later the Calcium phosphate crystals were added. The next morning cells were shocked for three minutes with 10% glycerol in a Tris buffered saline solution (TBS). The cells were harvested 48 h later by scraping with a rubber policeman.

Reporter gene assays

For the experiments involving the p18 KRAB-ER chimeras CAT reporter constructs were used, while pCMV-Luciferase served as an internal control. CAT activity was determined by the quantitative mixed-phase assay as described previously (35). The more recent experiments involving the Kox1, ZNF133 and ZNF140 KRAB ER chimeras the dual luciferase assay (Promega) was used according to manufacturers instructions with the PRL-SV40 serving as an internal control.

Results

To identify the most potent ERG-repressor, we separately co-transfected into CHO cells the constitutively active EREVITCAT promoter (36) and each of 7 different p18 KRAB repressors (Fig. 1). The basal activity of the EREVITCAT promoter in the absence of transfected reporter was established as a reference. Wild-type ER in the presence of ligand elicited a 2.2 fold increase compared to the no ligand control. The ERK repressor was the most effective repressing the basal activity of the promoter by 54% (Fig. 1). To examine repression in more detail we performed a dose response curve EREVITCAT promoter and the ERK and $\Delta A/B$ -ERK repressors. Clearly, ERK is the most potent of the two yielding 85% repression at the 100 ng dose (Fig. 2). While these data suggested that ERK would be a potent ERG-repressor, subsequent studies demonstrated that ERK would not provide a generally useful ERG-repressor. ERK failed to repress activity of the constitutively active 2ERETKCAT promoter (37) when wild-type ER was present, producing only a 38.5% inhibition of promoter activity at a 20:1 ratio of ERK:ER (Fig. 3).

Although the p18 repressors were effective under some conditions, repression was neither as universal nor as potent as desired. The KRAB domain can be subdivided into two sub domains the KRAB A and the KRAB B domain. The KRAB A domain is absolutely required for repression activity, while the KRAB B domain can further potentiate this action. The p18 KRAB domain only consists of the KRAB A domain. Earlier studies using GAL4 binding domain fusions compared p18 KRAB with other KRAB domains containing both the A and B sub domains (22), appeared to indicate that at higher levels of transfected KRAB the levels of repression converged. However, we could not exclude the possibility that the KRAB A domain by itself would be substantially weaker than the A + B domains together in a situation where the KRAB domain had to compete with the endogenous activation domains present in the ER to which it was fused. We therefore requested the cDNA clones of the Kox1 (or ZNF10, see ref. 25), ZNF133 and ZNF140 genes (24,26), all of which contain KRAB A+B

repression domains. Kox1 has been described most widely in literature. However, work by Vissing et al. (26) demonstrates that all three of these candidates should exhibit approximately equal potency as gene repressors. To exclude the possibility that lack of repression was due to an unusual interaction with the estrogen receptor, we also cloned GAL4-KRAB fusions and tested these on the G5-pGL3-Control vector in HepG2 cells (Fig. 4). The GAL4-Kox1 fusion potently repressed transcription. In this experiment the ZNF133 domain had approximately equal to slightly lower potency. We therefore are focusing our studies on the Kox1 variants of the various ERG repressors that we have recently cloned recently.

Our next experiments were then to establish whether the ER-Kox1 variant would also prove to be a potent transcription repressor (Fig. 5). In the first experiment we set out to establish a dose response curve for the KER10 variant. At 5 ng of transfected receptor we achieved 57% repression, with repression reaching a maximum of 71% at 40 ng of transfected receptor. At higher levels KER10 begins to affect the internal standard as well, which is reflected in the graph by the fact that the amount of repression appears to diminish, while in absolute terms repressor was more potent. The Kox1 receptors clearly are more potent at lower doses than the p18 KRAB constructs in CHO cells. This is particularly true since higher levels of expression are usually achieved in CHO cells from the CMV promoter compared to HepG2 cells.

To determine whether ablation of AF2 (as in KERFS10) or AF1 (as in $\Delta A/B$ -KER10) function, or the addition of a second KRAB domain (as in KERK10) could further potentiate repression, we compared the potency of a sub set of constructs at a single dose. While differences are small, the results show that both the deletion of AF1 and the addition of a second KRAB further potentiate repression (Fig. 6). The lack of potentiation by the ablation of AF2 function may result from the fact that this particular mutation causes the receptor to have a lowered affinity for its response element (32). The next obvious question then became whether the repressor would function in a ligand dependent fashion. HepG2 cells do not contain endogenous estrogen receptor. In the absence of transfected receptor the absence or presence of ligand (10^{-8} M Moxestrol, a slowly metabolized estrogen) has no influence on the level of activity of the promoter. Estrogen receptor robustly activated transcription in the presence of ligand alone. The KERK10 repressor potently repressed transcription in the presence of ligand, but only marginally in the absence of ligand (Fig. 7).

Finally, the KERK10 repressor can also potently repress both the ability of the wild-type ER to activate transcription from the EREs in the promoter and the constitutive activity of the promoter itself in the presence of wild type estrogen receptor. We compared the levels of repression in the presence and absence of 10 ng of cotransfected hER. The results demonstrate that at higher doses of KERK repression is hardly affected by the presence of estrogen receptor (Fig. 8). We would therefore expect the KERK10 to be an effective ERG-repressor in both the presence and absence of estrogen receptor.

Recommendations in Relation to the Statement of Work

The statement of work was prepared for a three year time frame with the first task to be completed in year one and task 2 within the first 18 months.

Task 1: Months 1-12: The preparation of potent repressors of estrogen regulated genes.

1. The use of the bacteriophage P22 challenge phage system to generate ER mutants with an enhanced affinity for the ERE. This task is essentially completed. We have identified ER-DBDs with >10 fold higher affinity for the ERE compared to the wild type ER DBD and full length receptors with up to 5-fold increase in affinity (TABLE 1). These mutants are also very effective in binding to a naturally occurring non consensus binding site, the ERE in the PS2 gene.
2. The insertion of the KRAB repressor domain at either the N- or C-terminus or both of an inactive estrogen receptor mutant. This task has been completed not only for the p18 KRAB domain but also for the Kox1, ZNF133 and ZNF140 KRAB domain. We will now focus our attention on the Kox1-ER chimeric repressors. We have not yet established the effectiveness of all the many Kox1-ER chimeras we constructed, but have shown that the sub set we tested are extremely efficient.
3. The replacement of the wild-type DNA binding domain of the KRAB repressor-ER construct with the DNA binding domain with the most strongly enhanced affinity for consensus and imperfect EREs. Our work in the P22 challenge phage system has demonstrated that there is no direct correlation between the potency of binding of the DBD mutants vs. The full length receptors. We have identified full-length receptors with an increased affinity and will construct KRAB chimeras with these. Interestingly we find that when fused to the KRAB domain these mutated DBDs can form effective repressors. While under certain circumstances it may be advantageous to have potent repressors that do not require ligand to induce the repressive phenotype, the use of full-length ER-KRAB chimeras may be warranted since recent reports have indicated that the ER not only acts directly by binding to the ERE, but can also act for instance through the AP1 (38) and the raloxifene response element (39), presumably via (an) intermediary protein(s). The A/B domain has been implicated in the interaction of ER with the c-Jun protein. It is not clear whether these are unique cases or whether this forms an emerging trend. It may well be that more promoters will be identified where ER acts through a similar mechanism, especially taking in mind that the presence of EREs has not been established in all estrogen regulated genes. In fact, our DBD-KRAB constructs may prove to be essential tools in identifying genes that are activated through those kind of mechanisms.

Task 2: Months 6-18: The Characterization of the ability of the ERG repressors to suppress transcription of synthetic reporter genes and of endogenous ER-regulated cellular genes.

1. Transient transfections to introduce the ERG repressors and reporter genes onto mammalian cell and into ER⁺ MCF7 and ER⁻ MDA-MB-231 breast cancer cells to determine their ability to repress transcription of synthetic reporter genes and of some endogenous genes including growth factor and growth factor receptor genes known to be under estrogen control. Our initial transient transfection studies using synthetic promoters have established the effectiveness of a sub set of the ERG repressors in HepG2 cells in both the absence and presence of ER. We are currently starting experiments in ER⁺ MCF7 and ER⁻ MDA-MB-231 breast cancer cells and expect to finish these studies within the projected time frame.
2. The construction and evaluation of the suppression by additional ERG-repressors. We have obtained the Kox1, ZNF133 and ZNF140 KRAB repressor domains and demonstrated that these are more potent than the p18 KRAB domain. We do not expect it to be necessary to construct further repressors containing the polycomb repressor (40).

CONCLUSIONS

The work carried out so far shows that KRAB-ER chimeras form extremely potent repressors that can effectively repress transcription from estrogen responsive genes, even in the presence of liganded wild-type estrogen receptor. While our initial constructs with the p18 KRAB were neither as potent nor as universal as hoped, the construction of the Kox1 KRAB chimeras has yielded powerful controllable ligand-dependent ERG-repressors.

Related work with the P22 challenge phage system has yielded estrogen DBD mutants that display an enhanced affinity for the estrogen response element. So far we have constructed DBD-KRAB chimeras with two of the most promising DBD mutants. In our initial test (Fig 9) these mutants proved to be very effective repressors of transcription. While the DBDs display up to a many fold higher affinity for the ERE, the full-length constructs have up to a 5-fold higher affinity. We will construct chimeric ER-KRAB constructs to test whether this raised affinity for the ERE will result in a more potent repressor.

REFERENCES

1. Herman, M. E., and Katzenellenbogen, B. S. (1994) Alterations in Transforming Growth Factor α and β Production and Cell Responsiveness During the Progression of MCF-7 Human Breast Cancer Cells to Estrogen-Autonomous Growth. *Cancer Res* 54, 5867-5874.
2. Nicholson, R. I., Gee, J. M. W., Francis, A. B., Manning, D. L., Wakeling, A. E., and Katzenellenbogen, B. S. (1995) Observations Arising from the Use of Pure Antioestrogens on Oestrogen-Responsive (MCF-7) and Oestrogen Growth-Independent (K3) Human Breast Cancer Cells. *Endocrine Related Cancer* 2, 1-7.
3. Arteaga, C. L., and Osborne, C. K. (1991) Growth Factors as Mediators of Estrogen/Antiestrogen Action in Human Breast Cancer Cells. In *Regulatory Mechanisms in Breast Cancer* (M. E. Lippman and R. B. Dickson, Eds.). Kluwer Academic Press, Boston, pp. 289.
4. Nicholson, R. I., McClelland, R. A., Gee, J. M. W., Manning, D. L., Cannon, P., Robertson, J. F. R., Ellis, I. O., and Blamey, R. W. (1994) Transforming Growth Factor α and Endocrine Sensitivity in Breast Cancer. *Cancer Res.* 54, 1684-1689.
5. Wiseman, L. R., Johnson, M. D., Wakeling, A. E., Lykkesfeldt, A. E., May, F. E. B., and Westley, B. R. (1994) Type-I IGF Receptor and Acquired Tamoxifen Resistance in Oestrogen-Responsive Human Breast Cancer Cells. *Eur. J. Cancer* 29A, 2256-2264.
6. Jordan, V. C. (1988) The Control of Hormone-Dependent Breast Cancer Growth--Are We Talking About Estrogen Alone? *Eur. J. Clin. Cancer Oncol.* 24, 1245-1248.
7. Dubik, D., and Shiu, R. P. (1992) Mechanism of Estrogen Activation of C-Myc Oncogene Expression. *Oncogene* 7, 1587-1594.
8. Dubik, D., and Shiu, R. P. (1988) Transcriptional Regulation of C-Myc Oncogene Expression by Estrogen in Hormone-Responsive Human Breast Cancer Cells. *J. Biol. Chem.* 263, 12705-12708.
9. Foster, J. S., and Wimalasena, J. (1996) Estrogen Regulates Activity of Cyclin-Dependent Kinases and Retinoblastoma Protein Phosphorylation in Breast Cancer Cells. *Mol. Endocrinol.* 10, 488-498.
10. Sutherland, R. L., Watts, C. K. W., and Musgrove (1995) Cell Cycle Control by Steroid Hormones in Breast Cancer: Implications for Endocrine Resistance. *Endocrine Related Cancer* 2, 87-96.
11. Gillett, C., Fantl, V., Smith, R., Fisher, C., Bartek, J., Dickson, C., Barnes, D., and Peters, G. (1994) Amplification and Overexpression of Cyclin D1 in Breast Cancer Detected by Immunohistochemical Staining. *Cancer Res.* 54, 1812-1817.

12. Cho, H., Ng, P. A., and Katzenellenbogen, B. S. (1991) Differential Regulation of Gene Expression by Estrogen in Estrogen Growth-Independent and -Dependent MCF-7 Human Breast Cancer Cell Sublines. *Mol. Endocrinol.* 5, 1323-1330.
13. Stewart, A. J., Johnson, M. D., May, F. E. B., and Westly, B. R. (1990) Role of Insulin-Like Growth Factors and the Type I IGF Receptor in the Estrogen-Stimulated Proliferation of Human Breast Cancer. *J. Biol. Chem.* 265, 72-78.
14. Stewart, A. J., Westley, B. R., and May, F. E. B. (1992) Modulation of the Proliferative Response of Breast Cancer Cells to Growth Factors by Oestrogen. *Br. J. Cancer* 66, 640-648.
15. Cullen, K. J., Yee, D., Sly, W. S., Perdue, J., Hampton, B., Lippman, M. E., and Rosen, N. (1990) Insulin-like Growth Factor Receptor Expression and Function in Human Breast Cancer. *Cancer Res.* 50, 48-53.
16. Dickson, R. B., and Lippman, M. E. (1987) Estrogenic Regulation of Growth and Polypeptide Growth Factor Secretion in Human Breast Carcinoma. *Endocr. Rev.* 8, 29-43.
17. Katzenellenbogen, B. S. (1991) Antiestrogen Resistance: Mechanisms by Which Breast Cancer Cells Undermine the Effectiveness of Endocrine Therapy. *J. Natl. Cancer Inst.* 83, 1434-1435.
18. Clarke, R., Br  nner, N., Katzenellenbogen, B. S., Thompson, E. W., Norman, M. J., Koppi, C., Soonmyoung, P., Lippman, M. E., and Dickson, R. B. (1989) Progression of Human Breast Cancer Cells from Hormone-Dependent to Hormone-Independent Growth Both *in vitro* and *in vivo*. *Proc Natl Acad Sci USA* 86, 3649-3653.
19. Clarke, R., Dickson, R. B., and Lippman, M. E. (1992) Hormonal Aspects of Breast Cancer: Growth Factors, Drugs and Stromal Interactions. *Critical Reviews in Oncology and Hematology* 12, 1-23.
20. Knabbe, C., Lippman, M. E., Wakefield, L. M., Flanders, K. C., Derynck, R., Dickson, R. B., and Kasid, A. (1987) Evidence that Transforming Growth Factor b is a Hormonally Regulated Negative Growth Factor in Human Breast Cancer Cells. *Cell* 48, 417-428.
21. Knabbe, C., Zugmaier, G., Schmahl, M., Dietel, M., Lippman, M. E., and Dickson, R. B. (1991) Induction Of Transforming Growth Factor B by the Antiestrogens Droloxifene, Tamoxifen, and Toremifene in MCF-7 Cells. *Am. J. Clin. Oncol.* 14(supp. 2), S15-S20.
22. Pengue, G., Calabro, V., Bartoli, P.C. Pagliuca, A., and Lania, L. (1994) Repression of Transcriptional Activity At A Disteance by the Evolutionarily Conserved Krab Domain Present in a Subfamily of Zinc Finger Proteins. *Nucl. Acids Res.* 22, 2908-2914.

23. Witzgall, R., O'Leary, E., Leaf, A., Onaldi, D., and Bonventre, J.V. (1994) The Kruppel-associated Box-A (LRAB-A) Domain of Zinc Finger Proteins Mediates Transcriptional Repression. *Proc. Nat'l. Acad. Sci. USA* 91, 4514-4518.
24. Margolin, J.F., Friedman, J.R., Meyer, W., Vissing, H., Thiesen, H.-J., and Rauscher, F.J. III (1994) Kruppel-associated Boxes are Potent Transcriptional Repressor Domains. *Proc. Nat'l. Acad. Sci. USA* 91, 4509-4513.
25. Thiesen, H.-J. (1991) Multiple Genes Encoding Zinc Finger Domains Are expressed in Human T Cells. *The New Biologist* 2(4), 363-374
26. Vissing, H., Meyer, W.K.-H, Aagaard, L. Tommerup, N. and Thiesen, H.-J. (1995) Repression of transcriptional Activity by Heterologous KRAB Domains present in Zinc Finger Proteins. *FEBS Letters* 369, 153-157.
27. Pengue, G., Caputo A., Rossi, C., Barbanti-Brodano, G., and Lania, L. (1995) Transcriptional Silencing of Human Immunodeficiency Type I Long Terminal Repeat-Driven Gene Expression by the Kruppel Associated Box Repressor Domain Targeted to the Transactivating Response Element. *J. Virol.* 69, 6577-6580.
28. Pengue, G., and Lania, L. (1996) Kruppel-Associated Box Mediated Repression of Rna Polymerase II Promoters is Influenced by the Arrangment of Basal Promoter Elements. *Proc. Nat'l. Acad. Sci. USA* 92, 1015-1020.
29. Benson, N., Sugionon, P., Mendleman, L.V. and Youderian, P. (1986) General Selection for Specific DNA Binding Activities. *Genetics* 114,1-14.
30. Lebreton, B., Prasad, P., Jayaram, M., and Youderian, P. (1988) Mutations that Improve the Binding of Yeast FLP Recombinase to Its Substrate. *Genetics* 118, 393-400.
31. Wrenn, C. K., and Katzenellenbogen, B. S. (1993) Structure-Function Analysis of the Hormone Binding Domain of the Human Estrogen Receptor by Region-Specific Mutagenesis and Phenotypic Screening in Yeast. *J. Biol. Chem.* 268, 24089-24098.
32. Ince, B.A., Zhuang, Y., Wrenn, C.K., Shapiro, D.J., and Katzenellenbogen, B.S. (1993) Potent Dominant Negative Mutants of the Human Estrogen Receptor. *J. Biol. Chem.* 268, 14026-14032.
33. Mattick, S., Glenn, K., de Haan, G., and Shapiro, D.J. (1997) Analysis of Ligand Dependence and Hormone Response Element Synergy in Transcription by Estrogen Receptor. *J. Steroid Biochem. Molec. Biol.*, 60(5-6), 285-294.
34. Lillie, J.W., and Green, M.R. (1989) Transcription Activation by the Adenovirus E1a Protein. *Nature* 338, 39-44.

35. Zhuang, Y., Katzenellenbogen, B. S., and Shapiro, D. J. (1995) Estrogen Receptor Mutants Which Do Not Bind 17 β -Estradiol Dimerize and Bind to the Estrogen Response Element *in vivo*. *Mol. Endocrinol.* 9, 457-466.
36. Chang, T.C., Nardulli, A.M., Lew, D., and Shapiro, D.J. (1992) The Role of Estrogen Response Elements in Expression of the *Xenopus laevis* Vitellogenin B1 Gene. *Molec Endocrinol.* 6, 346-354.
37. Reese, J.R., and Katzenellenbogen, B.S. (1991) Differential DNA Binding Abilities of Estrogen Receptor Occupied with Two Classes of Antiestrogens: Studies Using Human Estrogen Receptor Over-Expressed in Mammalian Cells. *Nucleic Acids Res.* 19, 6595-6602
38. Webb, P., Lopez, G. N., Uht, R. M., and Kushner, P. J. (1995). Tamoxifen Activation of the Estrogen Receptor/AP-1 pathway: Potential Origin for the Cell-Specific Estrogen-Like Effects of Antiestrogens. *Mol. Endocrinol.* 9, 443-456.
39. Na N. Yang, Murali Venugopalan, Hardikar, S., and Glasbrook, A. (1996) Identification of an estrogen Response Element Activated by Metabolites of 17 β -Estradiol and Raloxifene. *Science* 273, 122-1225
40. Bunker, C.A., and Kingston, R.E. (1995) Transcriptional Repression By *Drosophila* And Mammalian Polycomb Group Proteins In Transfected Mammalian Cells. *Molec. Cell. Biol.* 14, 1721-1732.




Figure 1. Effect of various p18KRAB-ER chimeras on EREVITCAT in CHO cells
In this Experiment the various chimeras were transfected at a 10ng dose. Plasmid CMV Luciferase (50 ng) served as an internal control and EREVITCAT (2 g) served as the reporter.




Figure 2. Transcription Repression by p18KRAB-ER chimeras ERK and Δ A/B-ERK on EREVITCAT in CHO cells. In this experiment the chimeras were transfected at doses varying from 1-100ng to establish a dose response curve. The basal activity of the promoter is established by transfection of the reporter in the absence of Effector (bar marked Alone). ERK was a more potent repressor that Δ A/B-ERK and achieved 85% repression. Estrogen receptor (1 ng) in the absence and presence of 10^{-8} M Estradiol served as a positive control. Plasmid CMV Luciferase (50 ng) served as an internal control and EREVITCAT (2 μ g) as the reporter.




Figure 3. Repression of p18KRAB-ER chimera ERK on (ERE)₂-TK-CAT in the presence of ER in CHO cells. Increasing amounts of ERK were cotransfected with 1ng ER at ratios indicated. The basal activity of the promoter is established by transfection of the reporter in the absence of Effector (bar marked Alone). While the promoter activity is decreased in the presence of ERK, a maximum of only 38.5% inhibition is achieved. Plasmid CMV Luciferase (50 ng) served as an internal control and EREVITCAT (2 g) as the reporter.




Figure. 4. Effects of various Flag-GAL4 based Effectors on the 5'-G5-pGL3-Control reporter plasmid in HepG2 Cells. The basal activity of the SV40 promoter/enhancer was determined in the absence of effector (indicated by the white bar). Two amounts (50 and 800 ng) of the Flag-GAL4 binding domain served as a negative control, while 25 ng transfected Flag-GAL4-VP16 served as a positive control. 50-800ng of Flag-Gal4-Kox1 KRAB, (black bars) was transfected and repressed transcription up to 96% repression. The Flag-GAL4-ZNF133 KRAB fusion displayed similar repression as the of Flag-Gal4-Kox1 KRAB construct.




Figure 5. Repression by increasing amounts (0-160 ng) of the N-terminal Kox1 Krab ER chimera KER10 on the (ERE)₄-pGL3-Control plasmid in HepG2 Cells. A maximum of 73% inhibition was achieved at 40 ng transfected effector. Plasmid (ERE)₄-pGL3-Control (200 ng) served as reporter and PRL-SV40 (200 ng) as internal control.




Figure 6. Repression of transcription by various ER-Kox1 KRAB chimeras on (ERE)₄-pGL3-Control plasmid in HepG2 Cells. Chimeras KER10, KERFS10, ΔA/B-ERK10 and KERK10 were transfected at 40 ng dose. Ablation of AF2 did not result in more potent repression. However, both the addition of an extra KRAB domain or the deletion of AF1 increased repression. Plasmid (ERE)₄-pGL3-Control (200 ng) served as reporter and PRL-SV40 (200 ng) as internal control.




Figure 7. Ligand Inducible repression of Transcription by the ER-Kox1 KRAB chimeras KERK10 on the (ERE)₄-pGL3-Control plasmid in HepG2 Cells. In this experiment indicated amounts of hER and KERK10 were transfected in the absence or presence of ligand (10⁻⁸M Moxestrol). Activity of the promoter in the absence of ligand was not affected by hER, but was slightly repressed by KERK10. Ligand clearly induced repression. Plasmid (ERE)₄-pGL3-Control (200 ng) served as reporter and PRL-SV40 (200 ng) as internal control.




Figure 8. Repression by ER-Kox1 KRAB chimera KERK10 at varying amounts in the absence or presence of cotransfected hER on plasmid $(ERE)_4$ -pGL3-Control in HepG2 Cells. In this experiment indicated amounts of KERK10 were transfected in the absence or presence of hER (10 ng). At higher amounts of transfected KERK10 the curves converge indicating that repression potency of KERK10 is not affected by presence of hER. Plasmid $(ERE)_4$ -pGL3-Control (200 ng) served as reporter and PRL-SV40 (200 ng) as internal control.




Figure 9. Repression on plasmid $(ERE)_4$ -pGL3-Control in HepG2 cells by mutant DBD #15 when fused to the Kox1 KRAB domain. Even at low dose of transfected DNA mutant DBD #15 which has a 15-fold increased affinity for the ERE (see TABLE 1) is an effective repressor of transcription. Plasmid $(ERE)_4$ -pGL3-Control (200 ng) served as reporter and PRL-SV40 (200 ng) as internal control.

TABLE 1. Affinity for the estrogen response element displayed by mutant DBDs selected in the p22 Challenge phage system directly and when inserted into the full length estrogen receptor

Clone Number	Affinity Mutant DBDs (fold over WT)	Affinity Full Length Receptor Mutants (fold over WT)
17	15.0	1.0
42	15.0	0.1
15	13.0	0.5
22	7.0	2.0
40	0.5	5.0



DEPARTMENT OF THE ARMY
US ARMY MEDICAL RESEARCH AND MATERIEL COMMAND
504 SCOTT STREET
FORT DETRICK, MARYLAND 21702-5012

REPLY TO
ATTENTION OF:

MCMR-RMI-S (70-1y)

23 Aug 01

MEMORANDUM FOR Administrator, Defense Technical Information
Center (DTIC-OCA), 8725 John J. Kingman Road, Fort Belvoir,
VA 22060-6218


SUBJECT: Request Change in Distribution Statement

1. The U.S. Army Medical Research and Materiel Command has reexamined the need for the limitation assigned to the technical reports listed at enclosure. Request the limited distribution statement for these reports be changed to "Approved for public release; distribution unlimited." These reports should be released to the National Technical Information Service.

2. Point of contact for this request is Ms. Judy Pawlus at DSN 343-7322 or by e-mail at judy.pawlus@det.amedd.army.mil.

FOR THE COMMANDER:

Encl


PHYLLIS M. RINEHART
Deputy Chief of Staff for
Information Management

Reports to be Downgraded to Unlimited Distribution

ADB241560	ADB253628	ADB249654	ADB263448
ADB251657	ADB257757	ADB264967	ADB245021
ADB263525	ADB264736	ADB247697	ADB264544
ADB222448	ADB255427	ADB263453	ADB254454
ADB234468	ADB264757	ADB243646	
ADB249596	ADB232924	ADB263428	
ADB263270	ADB232927	ADB240500	
ADB231841	ADB245382	ADB253090	
ADB239007	ADB258158	ADB265236	
ADB263737	ADB264506	ADB264610	
ADB239263	ADB243027	ADB251613	
ADB251995	ADB233334	ADB237451	
ADB233106	ADB242926	ADB249671	
ADB262619	ADB262637	ADB262475	
ADB233111	ADB251649	ADB264579	
ADB240497	ADB264549	ADB244768	
ADB257618	ADB248354	ADB258553	
ADB240496	ADB258768	ADB244278	
ADB233747	ADB247842	ADB257305	
ADB240160	ADB264611	ADB245442	
ADB258646	ADB244931	ADB256780	
ADB264626	ADB263444	ADB264797	